

# ORDSM0021: Clinical Conscious Sedation

View Online



---

[1]

Ashley, P.F. et al. 2018. Sedation of children undergoing dental treatment. Cochrane Database of Systematic Reviews. (Dec. 2018).  
DOI:<https://doi.org/10.1002/14651858.CD003877.pub5>.

[2]

Coulthard, P. et al. 2011. Estimating the need for dental sedation. 1. The Indicator of Sedation Need (IOSN) – a novel assessment tool. British Dental Journal. 211, 5 (Sep. 2011), E10-E10. DOI:<https://doi.org/10.1038/sj.bdj.2011.725>.

[3]

Craig, D. and Boyle, C.A. 2017. Practical conscious sedation. Quintessence Publishing Co. Ltd.

[4]

Henthorn, K. and Dickinson, C. 2010. The use of flumazenil after midazolam induced conscious sedation. British Dental Journal. 209, (2010).

[5]

Meechan, J.G. et al. 1998. Pain and anxiety control for the conscious dental patient. Oxford University Press.

[6]

Standards for Conscious Sedation in the provision of Dental Care and Accreditation —  
Royal College of Surgeons:  
<https://www.rcseng.ac.uk/dental-faculties/fds/publications-guidelines/standards-for-conscious-sedation-in-the-provision-of-dental-care-and-accreditation/>.