

ORDSM0021: Clinical Conscious Sedation

[View Online](#)

Ashley, P. F., Chaudhary, M., & Lourenço-Matharu, L. (2018). Sedation of children undergoing dental treatment. Cochrane Database of Systematic Reviews. <https://doi.org/10.1002/14651858.CD003877.pub5>

Coulthard, P., Bridgman, C. M., Gough, L., Longman, L., Pretty, I. A., & Jenner, T. (2011). Estimating the need for dental sedation. 1. The Indicator of Sedation Need (IOSN) – a novel assessment tool. *British Dental Journal*, 211(5), E10–E10. <https://doi.org/10.1038/sj.bdj.2011.725>

Craig, D., & Boyle, C. A. (2017). Practical conscious sedation (N. H. F. Wilson & J. G. Meechan, Eds.; Second edition, Vol. 15). Quintessence Publishing Co. Ltd.

Henthorn, K., & Dickinson, C. (2010). The use of flumazenil after midazolam induced conscious sedation. *British Dental Journal*, 209. <https://www.nature.com/articles/sj.bdj.2010.1132.pdf>

Meechan, J. G., Robb, N. D., & Seymour, R. A. (1998). Pain and anxiety control for the conscious dental patient. Oxford University Press.

Standards for Conscious Sedation in the provision of Dental Care and Accreditation — Royal College of Surgeons. (n.d.). <https://www.rcseng.ac.uk/dental-faculties/fds/publications-guidelines/standards-for-conscious-sedation-in-the-provision-of-dental-care-and-accreditation/>