

ORDSM0021: Clinical Conscious Sedation

[View Online](#)

[1]

J. G. Meechan, N. D. Robb, and R. A. Seymour, Pain and anxiety control for the conscious dental patient. Oxford: Oxford University Press, 1998.

[2]

D. Craig and C. A. Boyle, Practical conscious sedation, Second edition., vol. 15. London: Quintessence Publishing Co. Ltd, 2017.

[3]

P. F. Ashley, M. Chaudhary, and L. Lourenço-Matharu, 'Sedation of children undergoing dental treatment', Cochrane Database of Systematic Reviews, Dec. 2018, doi: 10.1002/14651858.CD003877.pub5.

[4]

P. Coulthard, C. M. Bridgman, L. Gough, L. Longman, I. A. Pretty, and T. Jenner, 'Estimating the need for dental sedation. 1. The Indicator of Sedation Need (IOSN) – a novel assessment tool', British Dental Journal, vol. 211, no. 5, pp. E10–E10, Sep. 2011, doi: 10.1038/sj.bdj.2011.725.

[5]

K. Henthorn and C. Dickinson, 'The use of flumazenil after midazolam induced conscious sedation', British Dental Journal, vol. 209, 2010 [Online]. Available: <https://www.nature.com/articles/sj.bdj.2010.1132.pdf>

[6]

'Standards for Conscious Sedation in the provision of Dental Care and Accreditation — Royal College of Surgeons'. [Online]. Available:
<https://www.rcseng.ac.uk/dental-faculties/fds/publications-guidelines/standards-for-conscious-sedation-in-the-provision-of-dental-care-and-accreditation/>