

ORDSM0021: Clinical Conscious Sedation

[View Online](#)

1.

Meechan, J.G., Robb, N.D., Seymour, R.A.: Pain and anxiety control for the conscious dental patient. Oxford University Press, Oxford (1998).

2.

Craig, D., Boyle, C.A.: Practical conscious sedation. Quintessence Publishing Co. Ltd, London (2017).

3.

Ashley, P.F., Chaudhary, M., Lourenço-Matharu, L.: Sedation of children undergoing dental treatment. Cochrane Database of Systematic Reviews. (2018).
<https://doi.org/10.1002/14651858.CD003877.pub5>.

4.

Coulthard, P., Bridgman, C.M., Gough, L., Longman, L., Pretty, I.A., Jenner, T.: Estimating the need for dental sedation. 1. The Indicator of Sedation Need (IOSN) – a novel assessment tool. British Dental Journal. 211, E10-E10 (2011).
<https://doi.org/10.1038/sj.bdj.2011.725>.

5.

Henthorn, K., Dickinson, C.: The use of flumazenil after midazolam induced conscious sedation. British Dental Journal. 209, (2010).

6.

Standards for Conscious Sedation in the provision of Dental Care and Accreditation —
Royal College of Surgeons,
<https://www.rcseng.ac.uk/dental-faculties/fds/publications-guidelines/standards-for-conscious-sedation-in-the-provision-of-dental-care-and-accreditation/>.